

injuries and non-communicable diseases.⁴ Underlying the increasing prevalence of diabetes are complex genetic, environmental and lifestyle factors, including infant malnutrition, dietary changes and lack of physical activity.⁵⁻⁷

Our large cross-sectional survey in rural Bangladesh in 2016 estimated the prevalence of diabetes to be 8.9% and 11.4% among men and women aged ≥ 30 years, respectively (Fottrell E, Submitted, 2018). Further, approximately 17% of men and 23% of women were identified to have impaired fasting glucose or impaired glucose tolerance, collectively termed intermediate hyperglycaemia. These estimates are comparable with other estimates.⁸⁻¹¹ Despite the high levels of diabetes and intermediate hyperglycaemia, awareness and control of the condition is low. In our 2016 survey, we found that only 25% of diabetics were aware of their status, women with diabetes were 37% less likely than men to know that they were diabetic and, even among known diabetics, 75% had suboptimal control of the condition (Fottrell E, Submitted, 2018). The Bangladesh Demographic and Health Survey also found low levels of awareness and control among a diabetic subsample,⁸ though general knowledge of diabetes prevention and control in the population was not investigated.

Reasons for low levels of awareness of one's diabetic status and control of it include a lack of widespread public health and awareness campaigns, inadequate health services for diagnosing and treating diabetes effectively, people not being able to afford regular healthcare and treatment, and communicable diseases remaining a priority for Bangladeshi public health programmes.¹² A study by Islam *et al* looking at knowledge of diabetes and glycaemic control among patients with diabetes in urban Dhaka, the capital, found that 46% of patients in the study had good, 38% moderate and 17% poor knowledge on diabetes.¹³ In a separate study in rural Bangladesh, knowledge that diabetes can cause eye disease and can be controlled by regular exercise was found to be higher among men.¹⁴ Unsurprisingly, knowledge of diabetes prevention, control, consequences and risk factors is significantly associated with higher education, higher monthly income, family history of diabetes and a longer duration of a diagnosis of diabetes.^{13 14}

In the current study, we describe diabetes knowledge and care practices among a large rural Bangladeshi population, measured as part of a baseline survey for a cluster randomised controlled trial for diabetes prevention and control led by the Diabetic Association of Bangladesh and University College London.¹⁵

METHODS

The study was conducted in Faridpur district and included 96 rural villages in four upazillas—Nagarkanda, Boalmari, Saltha and Madhukhali—covering a population of approximately 125 000 adults aged ≥ 30 years. Primary to

tertiary levels of healthcare are provided in Faridpur, but distance, long travelling time and a shortage of facilities, trained healthcare providers and medicines are ongoing challenges affecting access and quality of care.

The study population includes male and non-pregnant female permanent residents of the 96 villages aged ≥ 30 years. Someone was considered a permanent resident of a village if they normally live in that village. The study team conducted a census of all households and eligible residents to create a sampling frame from which a sample of 143 adults aged ≥ 30 years in each village was selected using multistage random sampling from a purpose-made sampling frame of all eligible individuals. In the first stage, 143 households with at least one eligible adult resident was selected using probability proportional to size sampling. At the next stage, a single eligible adult was selected for inclusion in the survey using simple random sampling. The sample size was determined by trial requirements described elsewhere¹⁵ but allows estimation of the true population prevalence of intermediate hyperglycaemia and diabetes mellitus with 99% confidence and an accuracy of between 1% and 2%.

DATA COLLECTION

Data were collected by 16 male and 16 female fieldworkers who were recruited locally and received approximately 1 month's training on survey methods. Male and female fieldworker pairs were supervised by one of four field supervisors who would spend at least half a day observing and verifying data within each pair at least every two days.

Field testing centres were established for the purposes of the study and were at a central, convenient location in each village. All consenting sampled individuals were requested to attend these centres on the morning of a specified day following an overnight fast for a range of physical measurements, including blood glucose. Blood glucose was measured using the One Touch Ultra Glucometer (Lifescan, Milpitas, California, USA) in whole blood obtained by finger prick from capillaries in the middle or ring finger after an overnight fast. All individuals then received a 75 g glucose load dissolved in 250 mL of water and had a repeat capillary blood test within 120 min (± 5 min) post ingestion to determine glucose tolerance status and differentiate between individuals with intermediate hyperglycaemia and those with diabetes according to WHO criteria¹⁶ or a prior medical diagnosis of diabetes.

Sociodemographic, lifestyle and behavioural data of all consenting individuals were collected through interview using a structured survey instrument. Questionnaire data were gathered using Samsung Galaxy Grand Prime smartphones using Open Data Kit Collect software. Collection of questionnaire data took place at the respondent's home before or after the physical measurements or at the testing centre at the time of physical measurement. Data were uploaded from mobile phones to the supervisors' laptop every two days and then transferred to a central database at the Faridpur field office for further

Malshah and colleagues¹⁰ being transferred on a weekly basis to the main project office in Dhaka.

Knowledge of diabetes was measured among all participants by asking a series of questions on (a) whether they were able to report any valid causes of diabetes, (b) whether they were able to report any valid symptoms of diabetes, (c) whether they were able to report any valid complications of diabetes, (d) whether they were able to report any valid ways to prevent diabetes and (e) whether they were able to report any valid ways to control diabetes if one has it. All of these questions were open-ended and unprompted, with fieldworkers ticking all valid responses from a checklist developed by the study team following review of the literature and discussion with diabetes specialists at the Bangladesh Institute of Research and Rehabilitation for Diabetes, Endocrine and Metabolic Disorders in Dhaka. The checklists were piloted in the study area and are available as online supplementary table 1. In addition, we gathered self-reported data on whether respondents had ever had their urine or blood tested for sugar. Finally, known diabetics were asked how often they had had their blood sugar checked since diagnosis.

A –

Descriptive analysis summarised levels of knowledge and practices by study population characteristics and diabetic status categorised as normoglycaemia (fasting glucose <6.1 mmol/L), intermediate hyperglycaemia (including impaired fasting glucose (fasting glucose \geq 6.1 mmol/L to <7.0 mmol/L) and 2-hour post ingestion of 75 g glucose load blood glucose <7.8 mmol/L) and impaired glucose tolerance (fasting glucose \geq 6.1 mmol/L to <7.0 mmol/L and 2-hour post ingestion of 75 g glucose load blood

t.0131Tj722/T1MC ETBT/T12 1 Tf10 0 0 10 191.31- 1 Tf10 0 0 3



Total			
--------------	--	--	--

Table 2 Adjusted ORs showing associations between study population characteristics and ability to correctly report valid answers within five domains of diabetes knowledge

		Ability to report valid answers within each of the following domains of knowledge of diabetes				
		Causes	Symptoms	Complications	Prevention	Control
		AOR 95% CI	AOR 95% CI	AOR 95% CI	AOR 95% CI	AOR 95% CI
Sex	Male					
	Female	1.02 (0.78 to 1.32)	1.13 (0.93 to 1.37)	1.05 (0.80 to 1.37)	0.99 (0.78 to 1.24)	0.96 (0.78 to 1.19)
Age (years)	30–39					
	40–49	1.02 (0.91 to 1.13)	1.03 (0.93 to 1.15)	1.08 (0.97 to 1.21)	1.01 (0.90 to 1.13)	1.03 (0.92 to 1.15)
	50–59	1.00 (0.88 to 1.13)	1.04 (0.91 to 1.19)	1.15 (0.99 to 1.34)	0.91 (0.78 to 1.05)	1.00 (0.87 to 1.15)
	60–69	0.84 (0.73 to 0.97)	0.73 (0.63 to 0.84)	0.94 (0.79 to 1.11)	0.71 (0.59 to 0.84)	0.66 (0.56 to 0.78)
	70+	0.45 (0.34 to 0.60)	0.42 (0.33 to 0.53)	0.51 (0.37 to 0.69)	0.47 (0.35 to 0.62)	0.41 (0.33 to 0.52)
Currently married	No					
	Yes	1.04 (0.87 to 1.25)	0.99 (0.85 to 1.16)	1.01 (0.85 to 1.20)	0.87 (0.73 to 1.04)	0.84 (0.71 to 1.00)
Education	No formal					
	Incomplete primary	1.22 (0.97 to 1.55)	1.67 (1.36 to 2.04)	1.61 (1.27 to 2.05)	1.47 (1.18 to 1.82)	1.91 (1.54 to 2.37)
	Completed at least primary	1.16 (0.84 to 1.60)	1.77 (1.34 to 2.33)	2.40 (1.78 to 3.23)	1.80 (1.36 to 2.39)	2.12 (1.56 to 2.88)
Literate	Illiterate					
	Literate	1.40 (1.07 to 1.83)	1.12 (0.88 to 1.42)	0.91 (0.71 to 1.18)	1.16 (0.89 to 1.50)	0.84 (0.66 to 1.08)
Occupation	Unemployed					
	Manual	1.12 (0.88 to 1.42)	1.17 (0.98 to 1.40)	1.15 (0.88 to 1.50)	1.11 (0.89 to 1.40)	1.18 (0.96 to 1.45)
	Professional	1.34 (1.08 to 1.66)	1.49 (1.21 to 1.83)	1.44 (1.13 to 1.85)	1.38 (1.11 to 1.71)	1.29 (1.02 to 1.64)
Wealth	Most poor					
	2	1.27 (1.07 to 1.52)	1.14 (0.97 to 1.33)	1.03 (0.85 to 1.24)	1.01 (0.85 to 1.19)	0.90 (0.72 to 1.12)
	3	1.27 (1.07 to 1.52)	1.41 (1.17 to 1.71)	1.20 (0.98 to 1.47)	1.29 (1.06 to 1.58)	1.38 (1.13 to 1.70)
	4	1.60 (1.29 to 1.98)	1.69 (1.37 to 2.09)	1.39 (1.08 to 1.78)	1.71 (1.35 to 2.16)	1.93 (1.59 to 2.35)
	Least poor	2.70 (2.14 to 3.41)	2.62 (2.01 to 3.41)	2.31 (1.79 to 2.99)	3.32 (2.55 to 4.31)	3.06 (2.33 to 4.02)
Religion	Other					
	Muslim	1.18 (0.82 to 1.70)	1.17 (0.85 to 1.60)	0.94 (0.66 to 1.33)	0.86 (0.63 to 1.18)	0.93 (0.62 to 1.40)
Diabetic status	Normoglycaemic					
	Intermediate hyperglycaemia	1.03 (0.89 to 1.18)	0.92 (0.79 to 1.07)	0.99 (0.86 to 1.14)	1.06 (0.93 to 1.20)	0.99 (0.87 to 1.14)
	Unknown diabetic	1.01 (0.82 to 1.23)	0.96 (0.79 to 1.18)	1.09 (0.87 to 1.36)	1.01 (0.82 to 1.25)	0.93 (0.75 to 1.14)
	Known diabetic	1.61 (1.23 to 2.09)	5.17 (3.41 to 7.82)	5.18 (3.75 to 7.14)	4.18 (3.04 to 5.74)	8.43 (4.83 to 14.71)

Results are adjusted for all covariates and for the stratified, clustered survey design. n for multivariate analysis=12045 due to missing occupation or diabetic status data for 95 individuals.

Unsurprisingly, almost all known diabetics reported ever having received a blood glucose test, and they were far more likely than any other group to have had either their blood or urine tested. Interestingly, unknown diabetics had significantly higher odds of blood and urine glucose testing relative to normoglycaemic individuals.

Among diabetic individuals who reported being awavdod

and diabetes knowledge was observed, with increasing knowledge among individuals from higher (less poor) wealth quintiles.

With regards to diabetic status, it is notable that there were no significant differences in knowledge between normoglycaemic individuals and those with intermediate hyperglycaemia or unknown diabetes. Known diabetics had significantly improved knowledge of diabetes, although levels remained low in absolute terms, with less than half knowing the cause of their disease.

The top ranking (frequency of 10% or more) reported causes, symptoms, complications, prevention and control strategies reported by respondents are summarised in [Box 1](#).

Only 14% of respondents reported ever having a blood glucose test, and fewer (5.2%) reported having a urine sugar test, with no differences observed between men and women ([table 3](#)). Increasing likelihood of blood or urine sugar testing was observed among older and more educated respondents. Observed crude associations between literacy and occupation became non-significant in multivariate analysis. Strong associations between wealth and ever having had a blood glucose test were observed, although even in the wealthiest quintile less than one-third of respondents had ever had a blood glucose test. The association between wealth and urine testing is less strong, with a significant association only being observed in the least poor group relative to the poorest.

Table 3 Frequency, crude and adjusted ORs and 95% CIs for blood glucose testing and urine glucose testing by sociodemographic characteristic

		Total	Ever blood glucose test			Ever urine glucose test			
		n	%	%	OR 95% CI	AOR 95% CI	%	OR 95% CI	AOR 95% CI
Sex	Male	5684	47.0	13.9			5.3		
	Female	6456	53.0	14.0	1.01 (0.90 to 1.13)	0.88 (0.68 to 1.15)	5.0	0.94 (0.79 to 1.11)	0.78 (0.55 to 1.11)
Age (years)	30–39	4108	33.6	9.8			3.5		
	40–49	3051	25.2	13.0	1.37 (1.20 to 1.57)	1.55 (1.33 to 1.81)	4.1	1.18 (0.94 to 1.49)	1.11 (0.86 to 1.45)
	50–59	2293	19.0	17.9	2.02 (1.72 to 2.37)	2.57 (2.13 to 3.08)	6.8	1.99 (1.57 to 2.52)	1.93 (1.45 to 2.56)
	60–69	1917	15.9	18.7	2.13 (1.80 to 2.51)	2.59 (2.09 to 3.22)	7.9	2.34 (1.88 to 2.90)	2.12 (1.60 to 2.81)
	70+	771	6.3	16.5	1.82 (1.46 to 2.27)	1.80 (1.32 to 2.45)	6.5	1.89 (1.35 to 2.66)	1.54 (0.99 to 2.41)
Currently married	No	1508	12.2	16.5			5.9		
	Yes	10632	87.8	13.6	0.80 (0.69 to 0.94)	0.76 (0.62 to 0.93)	5.1	0.85 (0.67 to 1.07)	0.88 (0.63 to 1.22)
Education	No formal	6057	49.5	9.4			3.6		
	Incomplete primary	2777	23.3	12.8	1.42 (1.21 to 1.66)	1.61 (1.26 to 2.05)	4.8	1.35 (1.07 to 1.71)	1.69 (1.21 to 2.36)
	Completed at least primary	3306	27.2	23.2	2.90 (2.47 to 3.41)	2.50 (1.86 to 3.35)	8.4	2.46 (1.97 to 3.07)	2.20 (1.39 to 3.48)
Literate	Illiterate	7475	61.4	10.0			3.8		
	Literate	4665	38.6	20.4	2.31 (2.03 to 2.63)	1.15 (0.90 to 1.47)	7.3	1.99 (1.66 to 2.39)	0.98 (0.67 to 1.45)
Occupation	Unemployed	6703	55.1	15.3			5.7		
	Manual	4034	33.3	9.1	0.55 (0.47 to 0.65)	0.58 (0.43 to 0.78)	3.1	0.53 (0.43 to 0.66)	0.52 (0.36 to 0.75)
	Professional	1401	11.6	21.7	1.54 (1.31 to 1.80)	0.87 (0.64 to 1.19)	8.4	1.51 (1.17 to 1.95)	0.89 (0.60 to 1.32)
Wealth	Most poor	2431	19.7	7.8			3.3		
	2	2445	20.4	7.4	0.94 (0.71 to 1.24)	0.84 (0.62 to 1.14)	3.4	1.04 (0.70 to 1.54)	0.97 (0.66 to 1.42)
	3	2441	20.2	11.1	1.47 (1.18 to 1.82)	1.32 (1.05 to 1.64)	4.4	1.37 (0.98 to 1.92)	1.21 (0.88 to 1.66)
	4	2403	19.6	14.4	1.97 (1.59 to 2.45)	1.60 (1.27 to 2.01)	4.8	1.50 (1.10 to 2.04)	1.16 (0.86 to 1.57)
	Least poor	2420	20.1	29.2	4.86 (3.95 to 5.98)	2.91 (2.32 to 3.66)	9.9	3.23 (2.36 to 4.42)	1.53 (1.09 to 2.13)
Religion	Other	1140	9.2	20.3			9.0		
	Muslim	11000	90.8	13.3	0.60 (0.47 to 0.77)	0.78			

study. Our questionnaire gathered data on unprompted knowledge of diabetes, which more realistically tests our respondents' ability to spontaneously recall and report their understanding of diabetes than relying on recognition of cases, symptoms, prevention and control when presented with a list of options or yes/no response categories. This approach is also more likely to reduce reporting bias.

our study is perhaps not surprising and supports other findings in both high-income and low-income settings.^{14 19 22-24} Health inequalities are well described in Bangladesh^{25 26} and the majority of the predicted rising burden of diabetes in Bangladesh is expected to occur in the low and middle socioeconomic groups where, as we have shown, knowledge of diabetes is poor and opportunities and ability to act to prevent and control the disease may be limited.^{18 19}

Current health services and preventative strategies in Bangladesh are inadequately prepared to address the challenges of high disease prevalence, looming increases in prevalence and low levels of knowledge, particularly in rural areas.^{18 21 27} In line with recommendations of the Global Strategy for the Prevention and Control of Non Communicable Diseases, there is a need for large-scale awareness intervention programmes that target not only high-risk individuals, but whole populations.^{28 29} Raising knowledge and awareness of diabetes in the wider population is necessary, not least because previous studies have shown that, in the absence of universal health coverage and access to professional services, the most common source of information on prevention and care seeking for diabetes are family members, friends and neighbours.²¹ The fact that we have observed even a small degree of knowledge within our study population may therefore be seen as a positive base on which to build and increase the spread of knowledge. Mass media health promotion campaigns and opportunities created by the widespread ownership of mobile phones may create opportunities for this. It is important, however, that any such strategies are tailored to the context and literacy of adult rural Bangladeshi populations and inclusive of those with lower education and wealth. Lessons may also be learnt from recent population health gains in Bangladesh, such as progress in maternal and child health, and group and individual knowledge, awareness and behaviour change interventions that have shown success in relation to other health outcomes. There is a need for any such innovations and interventions to be robustly evaluated and evidence-based before scale-up.

CONCLUSION

Our cross-sectional survey in a large rural population shows that knowledge on the causes, symptoms, consequences, prevention and control of diabetes is limited in rural Bangladesh. A minority of individuals with diabetes are aware of their status and even then do not appear to monitor their blood glucose levels on a regular basis. These findings may be reflective of low levels of health literacy and inadequate availability and access to health services and information in rural areas. Community-based interventions that promote knowledge and understanding of diabetes are needed and these should capitalise on existing knowledge and prevailing lay networks of as. Comclgxiswaviour ntione hy and p63 0 08jTw Tflxistss progress in r8-T10 1 Tf0 -ða T1J0.ss inniq1Dw basis. These/oK/icac//Tiw .e Lofrom recent population health gains ij08sfJMtion healeyk

